Carrey Islet Bird Checklist (Off Koumac NW GT) New Caledonia

20 23 00s 163 57 00e

Compiled by Michael K. Tarburton, Pacific Adventist University, PNG.

swiftlet@optusnet.com.au

[You are welcome to communicate, just re-type above address into your e-mail program]

| # | Common Name | Scientific Name | Ecol. Status | Abundance | References |
|----|--------------|---------------------------|--------------|---|------------|
| 1. | Brown Booby | Sula leucogasta plotus | Res bre | Br 1978, 200 birds incl 29 Pr on 21 Nov 2005, | 1, |
| 2. | Masked Booby | Sula dactylatra personata | Res bre | 1 pr with yng, 26 Aug 2005, | 1, |
| 3. | Sanderling | Calidris a. alba | Sum mig | 2 obs 26 Aug & 23 Nov 2005, | 1, |
| 4. | Fairy Tern | Sternula nereis exsul | Res | 3 pair 28 Aug 2005, | 1, |

<u>KEY</u>

Ecological status

End = Endemic to Island unless otherwise stated, **Res bre** = Resident breeder, **Res** = resident ie breeding not proven or not likely. For seabirds it means they have or should be found in that area: note other refs. **Vag** = Vagrant, **Res/mig** = Part of population migratory, **Sum mig** = Summer migrant, **Win mig** = Winter migrant, **Spr/Aut mig** = Passage migrant in Spring and Autumn, **Intro** = Introduced.

Abundance

 $\mathbf{R} = \text{rare}, \mathbf{S} = \text{scarce}, \mathbf{UC} = \text{uncommon}, \mathbf{MC} = \text{moderately common}, \mathbf{LC} = \text{locally common}, \mathbf{VC} = \text{very common}, \mathbf{A} = \text{abundant}, \mathbf{P} = \text{present}.$

Colour coding applies only to one species on a particular list. eg. An abundance comment in red for the Brown Booby refers to the Ref # in Red for that species only. **Alternate Island Names & alternate common Bird Names are recorded in two separate spreadsheet files.**

REFERENCES

 Barre, N. Baudat-Francheschi, J. Spaggiari, J. Chartendrault, V. Bachy, P. Desmooulins, F. Guhring, J. 2007. Second complement á la liste des oiseaux de Nouvelle-Calédonie. *Alauda* 75(2), 129-144.