Menyuru (Menyuro) Islet Bird Checklist

(Eastern Lagoon) New Caledonia

21 44 48s 166 30 53e

Compiled by Michael K. Tarburton, Pacific Adventist University, PNG.

swiftlet@optusnet.com.au

[You are welcome to communicate, just re-type above address into your e-mail program]

#	Common Name	Scientific Name	Ecol. Status	Abundance	References
1.	Fairy Tern	Sternula nereis exsul	Res bre	11 pr breeding 2008,	1

KEY

Ecological status

End = Endemic to Island unless otherwise stated, **Res bre** = Resident breeder, **Res** = resident ie breeding not proven or not likely. For seabirds it means they have or should be found in that area: note other refs. **Vag** = Vagrant, **Res/mig** = Part of population migratory, **Sum mig** = Summer migrant, **Win mig** = Winter migrant, **Spr/Aut mig** = Passage migrant in Spring and Autumn, **Intro** = Introduced.

Abundance

R = rare, S = scarce, UC = uncommon, MC = moderately common, LC = locally common, C = common, VC = very common, A = abundant, P = present.

Colour coding applies only to one species on a particular list. eg. An abundance comment in red for the Brown Booby refers to the Ref # in Red for that species only.

Alternate Island Names & alternate common Bird Names are recorded in two separate spreadsheet files.

REFERENCES

1. Barré, N., Baling, M., Baillon, N., Le Bouteiller, A., Bachy, P., Chartendrault, V. & Spaggiari, J. 2012. Survey of Fairy Terns *Sterna nereis exsul* in New Caledonia. *Marine Ornithology* 40, 31-38.