Ndie Islet Bird Checklist (Sthn Lagoon not one in Dumbea Bay,) Off New Caledonia

22 31 11s 167 11 53e

Compiled by Michael K. Tarburton, Pacific Adventist University, PNG.

swiftlet@optusnet.com.au

[You are welcome to communicate, just re-type above address into your e-mail program]

#	Common Name	Scientific Name	Ecol. Status	Abundance	References
1.	Wedge-tailed Shearwater	Ardenna pacificus	Res bre	1,000-2,000 br prs 1993-99,	2,
2.	Eastern Osprey	Pandion cristatus	Res bre	1 nest, presume 1 pr,	1,
3.	Brown Noddy	Anous stolidus	Res bre	1-5 pr bred 1993-99,	2,
4.	Black Noddy	Anous minutus	Res bre	5,001-10,000 pr bred 1993-99,	2,

KEY

Ecological status

End = Endemic to Island unless otherwise stated, **Res bre** = Resident breeder, **Res** = resident ie breeding not proven or not likely. For seabirds it means they have or should be found in that area: note other refs. **Vag** = Vagrant, **Res/mig** = Part of population migratory, **Sum mig** = Summer migrant, **Win mig** = Winter migrant, **Spr/Aut mig** = Passage migrant in Spring and Autumn, **Intro** = Introduced.

Abundance

 $\mathbf{R} = \text{rare}, \mathbf{S} = \text{scarce}, \mathbf{UC} = \text{uncommon}, \mathbf{MC} = \text{moderately common}, \mathbf{LC} = \text{locally common}, \mathbf{VC} = \text{very common}, \mathbf{A} = \text{abundant}, \mathbf{P} = \text{present}.$

Colour coding applies only to one species on a particular list. eg. An abundance comment in red for the Brown Booby refers to the Ref # in Red for that species only. Alternate Island Names & alternate common Bird Names are recorded in two separate spreadsheet files.

REFERENCES

1. Claude, MI. Panoraima photo on Google Earth showing Osprey nest. <u>www.panoramio.com/photo/62274708?source=wapi&referrer</u> downloaded 16.9.2012.

2. Benoit, M.P., & Bretagnolle, V. 2002. Seabirds of the Southern Lagoon of New Caledonia; Distribution, Abundance and Threats. Waterbirds 25(2), 202-213.