Ua Islet Bird Checklist (Sthn Lagoon) Off New Caledonia

22 42 31s 166 48 43e

Compiled by Michael K. Tarburton, Pacific Adventist University, PNG.

swiftlet@optusnet.com.au

[You are welcome to communicate, just re-type above address into your e-mail program]

•
3,
3,
2,
1,
2,
2,

KEY

Ecological status

End = Endemic to Island unless otherwise stated, **Res bre** = Resident breeder, **Res** = resident ie breeding not proven or not likely. For seabirds it means they have or should be found in that area: note other refs. **Vag** = Vagrant, **Res/mig** = Part of population migratory, **Sum mig** = Summer migrant, **Win mig** = Winter migrant, **Spr/Aut mig** = Passage migrant in Spring and Autumn, **Intro** = Introduced.

Abundance

 \mathbf{R} = rare, \mathbf{S} = scarce, \mathbf{UC} = uncommon, \mathbf{MC} = moderately common, \mathbf{LC} = locally common, \mathbf{C} = common, \mathbf{VC} = very common, \mathbf{A} = abundant, \mathbf{P} = present. Colour coding applies only to one species on a particular list. eg. An abundance comment in red for the Brown Booby refers to the Ref # in Red for that species only.

Alternate Island Names & alternate common Bird Names are recorded in two separate spreadsheet files.

REFERENCES

- 1. Barré, N., Baling, M., Baillon, N., Le Bouteiller, A., Bachy, P., Chartendrault, V. & Spaggiari, J. 2012. Survey of Fairy Terns Sterna nereis exsul in New Caledonia. Marine Ornithology 40, 31-38.
- 2. Google Earth Photo by Richard Chesher, at site of islet accessed 13.9.2012.
- 3. Benoit, M.P., & Bretagnolle, V. 2002. Seabirds of the Southern Lagoon of New Caledonia; Distribution, Abundance and Threats. Waterbirds 25(2), 202-213.