# Ugo Island Bird Checklist (off SW coast Grande Terre) New Caledonia

22 26 31s 166 55 44e

#### Compiled by Michael K. Tarburton, Pacific Adventist University, PNG.

# swiftlet@optusnet.com.au

#### [You are welcome to communicate, just re-type above address into your e-mail program]

<u>#</u>	Common Name	Scientific Name	Ecol. Status	Abundance	References
1.	Black-winged Petrel	Pterodroma nigripennis	Res bre	11-50 pr bred 1993-99,	2,
2.	Tahiti Petrel	Pseudobulweria rostrata	Res bre	1 pr bred 1993-1999,	2,
3.	Wedge-tailed Shearwater	Ardenna pacificus	Res bre	1,000-2,000 br prs 1993-99,	2,
4.	Crested Tern	Thalasseus bergii cristatus	Res	60 roosting & 11 en route there 25.6.2007	1,

## KEY

## **Ecological status**

End = Endemic to Island unless otherwise stated, **Res bre** = Resident breeder, **Res** = resident ie breeding not proven or not likely. For seabirds it means they have or should be found in that area: note other refs. Vag = Vagrant, **Res/mig** = Part of population migratory, **Sum mig** = Summer migrant, **Win mig** = Winter migrant, **Spr/Aut mig** = Passage migrant in Spring and Autumn, **Intro** = Introduced.

#### Abundance

 $\mathbf{R} = \text{rare}, \mathbf{S} = \text{scarce}, \mathbf{UC} = \text{uncommon}, \mathbf{MC} = \text{moderately common}, \mathbf{LC} = \text{locally common}, \mathbf{VC} = \text{very common}, \mathbf{A} = \text{abundant}, \mathbf{P} = \text{present}.$ 

Colour coding applies only to one species on a particular list. eg. An abundance comment in red for the Brown Booby refers to the Ref # in Red for that species only. Alternate Island Names & alternate common Bird Names are recorded in two separate spreadsheet files.

#### **REFERENCES**

1. Tarburton, M.K. 2007. Field records 2007.

2. Benoit, M.P., & Bretagnolle, V. 2002. Seabirds of the Southern Lagoon of New Caledonia; Distribution, Abundance and Threats. Waterbirds 25(2), 202-213.