Longan Systematic Bird Check List

(Ninigo Islands) Bismarck Archipelago, PNG.

1 13 12s 144 16 08e

Compiled by M.K. Tarburton, Pacific Adventist University, PNG.

swiftlet@optusnet.com.au

(for contact, re-type e-mail address)

#	Common Name	Scientific Name	Ecol. Status	Abundance	References
1.	Common Tern	Sterna hirundo	Sum mig	Small groups on lagoon side 1970.	2.
2.	Crested Tern	Thalasseus bergii	Res bre	Singles obs Aug 1970.	1,2,
3.	Black Noddy	Anous minutus	Res bre	A on lagoon side Aug 1970.	1,2,
4.	White Tern	Gygis alba	Res bre	C on adjacent lagoon Aug 1970.	1,2,
5.	Rainbow Lorikeet	Trichoglossus haematodus flavicans	Res bre	A 1970 obs flying from Menam,	2,
6.	Atoll Starling	Aplornis feadensis heureka	Res bre	Small numbers seen flying from Menam 1970,	2,

<u>KEY</u>

Ecological status

End = Endemic to Island unless otherwise stated, **Res bre** = Resident breeder, **Res** = resident ie breeding not proven or not likely. For seabirds it means they have or should be found in that area: note other refs. **Vag** = Vagrant, **Res/mig** = Part of population migratory, **Sum mig** = Summer migrant, **Win mig** = Winter migrant, **Spr/Aut mig** = Passage migrant in Spring and Autumn, **Intro** = Introduced.

Abundance

 $\mathbf{R} = \text{rare}, \mathbf{S} = \text{scarce}, \mathbf{UC} = \text{uncommon}, \mathbf{MC} = \text{moderately common}, \mathbf{LC} = \text{locally common}, \mathbf{C} = \text{common}, \mathbf{VC} = \text{very common}, \mathbf{A} = \text{abundant } \mathbf{P} = \text{present}.$

Colour coding applies only to one species on a particular list. eg. An abundance comment in red for the Brown Booby refers to the Ref # in Red for that species only. **Alternate Island Names & alternate common Bird Names are recorded in two separate spreadsheet files.**

REFERENCES

- 1. Bell, H.L. 1970. Ninigo & Hermit Islands, 8-15/8/1970. Papua New Guinea Bird Society Newsletter 57, 2-4.
- 2. Bell, H.L. 1975. Avifauna fo the Ninigo and Hermit Islands, New Guinea. Emu 75, 77-84.