# Mal Islet Systematic Bird Check List (Ninigo or L'Exchequier Islands) Bismarck Archipelago, PNG. 1 24 02s 144 10 29e

Compiled by M.K. Tarburton, Pacific Adventist University, PNG.

swiftlet@optusnet.com.au

[To communicate: please re-type e-mail address]

| #  | Common Name    | Scientific Name             | Ecol. Status | Abundance                          | References |
|----|----------------|-----------------------------|--------------|------------------------------------|------------|
| 1. | Black Myzomela | Myzomela negrita ernstmayri | Res bre      | Type coll. by Mayr 10-11 Nov 1928, | 1,         |
| 2. | Atoll Starling | Aplornis feadensis heureka  | Res bre      | Type coll. by Mayr 10-11 Nov 1928, | 1,         |

## <u>KEY</u>

### **Ecological status**

End = Endemic to Island unless otherwise stated, **Res bre** = Resident breeder, **Res** = resident ie breeding not proven or not likely. For seabirds it means they have or should be found in that area: note other refs. Vag = Vagrant, **Res/mig** = Part of population migratory, **Sum mig** = Summer migrant, **Win mig** = Winter migrant, **Spr/Aut mig** = Passage migrant in Spring and Autumn, **Intro** = Introduced.

#### Abundance

 $\mathbf{R}$  = rare,  $\mathbf{S}$  = scarce,  $\mathbf{UC}$  = uncommon,  $\mathbf{MC}$  = moderately common,  $\mathbf{LC}$  = locally common,  $\mathbf{C}$  = common,  $\mathbf{VC}$  = very common,  $\mathbf{A}$  = abundant  $\mathbf{P}$  = present.

Colour coding applies only to one species on a particular list. eg. An abundance comment in red for the Brown Booby refers to the Ref # in Red for that species only. Alternate Island Names & alternate common Bird Names are recorded in two separate spreadsheet files.

#### **REFERENCES**

1. Bell, H.L. 1975. Avifauna of the Ninigo and Hermit Islands, New Guinea. Emu 75, 77-84. Citing: Mayr, E. 1929, Neuguinea-Reise (Briefe), Ornithol Mber. 37, 62-64.