Haycock Stack #2 Bird Checklist

Second last of four stacks North of Haycock Isl., Solomon Islands

7 27 59.09s 157 52 31.58e

Compiled by Michael K. Tarburton, Pacific Adventist University, PNG.

swiftlet@optusnet.com.au

[To communicate: please re-type e-mail address]

| # | Common Name | Scientific Name | Ecol. Status | Abundance | References |
|----------|--|--|--------------|--|------------|
| | | | | | |
| 1. 2. | Black-naped Tern Island Imperial Pigeon | Sterna sumatrana sumatrana Ducula pistrinaria | Res Res | 2 present 21.12.2005, 5 birds 21.12.05, | 1, 1, |

KEY

Ecological status

End = Endemic to Island unless otherwise stated, **Res bre** = Resident breeder, **Res** = resident ie breeding not proven or not likely. For seabirds it means they have or should be found in that area: note other refs. **Vag** = Vagrant, **Res/mig** = Part of population migratory, **Sum mig** = Summer migrant, **Win mig** = Winter migrant, **Spr/Aut mig** = Passage migrant in Spring and Autumn, **Intro** = Introduced.

Abundance

R = rare, S = scarce, UC = uncommon, MC = moderately common, LC = locally common, C = common, VC = very common, A = abundant P = present.

Colour coding applies only to one species on a particular list. eg. An abundance comment in red for the Brown Booby refers to the Ref # in Red for that species only.

Alternate Island Names & alternate common Bird Names are recorded in two separate spreadsheet files.

REFERENCES

1. Tarburton, M.K. 2005. Personal Field Notes.