# Fenualoa (Nibarga or Nibanga Temau) Island Bird Checklist Reef Group, Santa Cruz Isles, Solomon Islands

### 10 14 14.61s 166 18 33.06e

Compiled by Michael K. Tarburton, Pacific Adventist University, PNG.

swiftlet@optusnet.com.au

[To communicate: please re-type e-mail address]

#	Common Name	Scientific Name	Ecol. Status	Abundance	References
1.	Eastern Reef Egret	Egretta sacra	Res	Р,	1,
2.	Red-bellied Fruit-Dove	Ptilinopus greyi	Res bre	Coll. 1920's	4,
3.	Pacific Emerald Dove	Chalcophaps longirostris sandwichensis	Res bre	Р,	1,
4.	Pacific Imperial Pigeon	Ducula pacifica	Res bre	Р,	1,
5.	Long-tailed Koel	Eudynamys taitensis	Win mig	M coll. Oct 1926.	1,2,
6.	White-rumped Swiftlet	Aerodramus spodiopygius leucopygia/des	olataRes bre	P, 2M skins AMNH,	1,
7.	Collared Kingfisher	Halcyon chloris brachyura	Res bre	Р,	1,
8.	Cardinal Myzomela	Myzomela cardinalis sanctacrucis	Res bre	P, Spec. AMNH	1,3,

## KEY

#### **Ecological status**

End = Endemic to Island unless otherwise stated, **Res bre** = Resident breeder, **Res** = resident ie breeding not proven or not likely. For seabirds it means they have or should be found in that area: note other refs. **Vag** = Vagrant, **Res/mig** = Part of population migratory, **Sum mig** = Summer migrant, **Win mig** = Winter migrant, **Spr/Aut mig** = Passage migrant in Spring and Autumn, **Intro** = Introduced.

#### Abundance

 $\mathbf{R} = \text{rare}, \mathbf{S} = \text{scarce}, \mathbf{UC} = \text{uncommon}, \mathbf{MC} = \text{moderately common}, \mathbf{LC} = \text{locally common}, \mathbf{C} = \text{common}, \mathbf{VC} = \text{very common}, \mathbf{A} = \text{abundant}, \mathbf{P} = \text{present}.$ 

Colour coding applies only to one species on a particular list. eg. An abundance comment in red for the Brown Booby refers to the Ref # in Red for that species only. Alternate Island Names & alternate common Bird Names are recorded in two separate spreadsheet files.

#### **REFERENCES**

- 1. Mayr E. 1945. Birds of the South-West Pacific. Wheldon Press, New York.
- 2. Bogert, C. 1937. Birds Collected during the Whitney South Sea Expedition. 34. The distribution and migration of the Long-tailed Cuckoo (*Urodynamis taitensis* Sparrman). *Am. Mus. Novit.* **933**: 1-12.
- 3. Mayr, E. 1932. Birds collected during the Whitney South Sea Expedition 18. Notes on Meliphagidae from Polynesia and the Solomon Islands. Am. Mus. Novit. 516: 1-30.
- 4. Ripley, S.D. & H. Birkhead. 1942. Birds collected during the Whitney South Sea Expedition.51. On the fruit pigeons of the Ptilinopus purpuratus group. *Amer. Mus. Novit.* **1192**, 1-14.

5. Salomonsen, F. 1983. Revision of the Melanesian Swiftlets (*Apodes, Aves*) & their Conspecific Forms in the Indo-Australian and Polynesian Region. Noona Dan Papers No. 141.