

Bélep Island Bird Checklist

New Caledonia

Compiled by Michael K. Tarburton, Pacific Adventist University, PNG.

swiftlet@optusnet.com.au

| # | Common Name | Scientific Name | Ecol. Status | Abundance | References |
|----|---------------------|--------------------------------|--------------|--|------------|
| 1. | Tahiti Petrel | <i>Pseudobulweria rostrata</i> | Res | Found Breeding 1988-2000, | 1, |
| 2. | Australasian Gannet | <i>Sula serrator</i> | Vag | Bird banded at Philip Isl found at Belep Jan 1996, | 1, |
| 3. | Masked Lapwing | <i>Vanellus miles</i> | Self Intro | Obs Oct 1995 & copulation obs Oct 1996, | 1, |
| 4. | Grey Plover | <i>Pluvialis squatarola</i> | Sum mig | Obs at Koumac, | 1, |

KEY

Ecological status

End = Endemic to Island unless otherwise stated, **Res bre** = Resident breeder, **Res** = resident ie breeding not proven or not likely. For seabirds it means they have or should be found in that area: note other refs. **Vag** = Vagrant, **Res/mig** = Part of population migratory, **Sum mig** = Summer migrant, **Win mig** = Winter migrant, **Spr/Aut mig** = Passage migrant in Spring and Autumn, **Intro** = Introduced.

Abundance

R = rare, **S** = scarce, **UC** = uncommon, **MC** = moderately common, **LC** = locally common, **C** = common, **VC** = very common, **A** = abundant, **P** = present.

Colour coding applies only to one species on a particular list. eg. An abundance comment in **red** for the Brown Booby refers to the Ref # in **Red** for that species only.

Alternate Island Names & alternate common Bird Names are recorded in two separate spreadsheet files.

REFERENCES

1. Barré, N. & Dutson, G. 2000. Liste commentée des oiseaux de Nouvelle-Calédonie. *Alauda* 3: suppl. 1-48.